

UTILIZATION MANAGEMENT MEDICAL POLICY

POLICY: Oncology (Injectable – HER2 Antagonist) – Trastuzumab Products Utilization Management Medical Policy

- Herceptin[®] (trastuzumab intravenous infusion – Genentech)
- Hercessi[™] (trastuzumab-strf intravenous infusion – Accord BioPharma)
- Herzuma[®] (trastuzumab-pkrb intravenous infusion – Celltrion)
- Kanjinti[™] (trastuzumab-anns intravenous infusion – Amgen)
- Ogivri[®] (trastuzumab-dkst intravenous infusion – Mylan)
- Ontruzant[®] (trastuzumab-dttb intravenous infusion – Merck)
- Trazimera[™] (trastuzumab-qyyp intravenous infusion – Pfizer)

REVIEW DATE: 06/24/2026

OVERVIEW

Trastuzumab products are human epidermal growth factor receptor 2 (HER2)/neu receptor antagonists indicated for the following uses:¹

- **Breast cancer, adjuvant treatment** of HER2-overexpressing node positive or node negative disease (estrogen receptor[ER]/progesterone receptor [PR] negative or with one high risk feature) 1) as part of a treatment regimen consisting of doxorubicin, cyclophosphamide, and either paclitaxel or docetaxel; 2) as part of a treatment regimen with docetaxel and carboplatin; or 3) as a single agent following multi-modality anthracycline based therapy.
- **Breast cancer, metastatic**, HER2-overexpressing, either in combination with paclitaxel for first-line treatment, or as a single agent in patients who have received one or more chemotherapy regimens for metastatic disease.
- **Gastric cancer or gastroesophageal (GE) junction adenocarcinoma, metastatic**, HER2-overexpressing disease, in combination with cisplatin and capecitabine or 5-fluorouracil (5-FU) who have not received prior treatment for metastatic disease.

Herzuma, Hercessi, Kanjinti, Ogivri, Ontruzant, and Trazimera are all approved biosimilars for Herceptin; all of the biosimilars have the same FDA-approved indications as Herceptin. For all indications, patients must be selected for therapy based on an FDA-approved companion diagnostic for trastuzumab. Tests are specific for breast cancer or gastric cancer.

Dosing Information

The approved dosing of trastuzumab for adjuvant treatment of breast cancer is given for a total of 52 weeks.¹ Initial dose is 4 mg/kg intravenously, then 2 mg/kg weekly for 12 weeks (with paclitaxel or docetaxel) or 18 weeks (with docetaxel/carboplatin). One week after the last weekly dose, trastuzumab 6 mg/kg is given every 3 weeks to complete a total of 52 weeks of therapy. Another dosing schedule is an initial dose of 8 mg/kg, then 6 mg/kg every 3 weeks for a total of 52 weeks of therapy. Extending adjuvant treatment beyond 1 year is not recommended. The approved dosing for metastatic breast cancer is trastuzumab (alone or in combination with paclitaxel) at an initial dose of 4 mg/kg given intravenously followed by weekly doses of 2 mg/kg until disease progression.¹ Many dosing schedules for trastuzumab are included in guidelines.² Alternate dosing will be assessed individually on a case-by-case basis.

The approved dose of trastuzumab given with chemotherapy in metastatic gastric cancer is an initial dose of 8 mg/kg intravenously followed by subsequent doses of 6 mg/kg every 3 weeks until progression.¹ Guidelines recommend either trastuzumab 8 mg/kg on Day 1 of Cycle 1 and then 6 mg/kg every 21 days

or trastuzumab 6 mg/kg on Day 1 of Cycle 1 and then 4 mg/kg every 14 days for first-line or second-line therapy (in combination with chemotherapy) for metastatic or locally advanced gastric, esophageal, or GE junction cancer.³⁻⁴

For colon cancer or rectal cancer, when used in combination with Perjeta® (pertuzumab intravenous infusion), trastuzumab is given as an 8 mg/kg infusion on Day 1 of Cycle 1 followed by 6 mg/kg every 21 days. When used in combination with lapatinib, trastuzumab is given as a 4 mg/kg infusion on Day 1 of Cycle 1, followed by 2 mg/kg weekly.⁵⁻⁶

For biliary tract cancer, appendiceal cancer, endometrial carcinoma and salivary gland tumors, in the clinical studies, trastuzumab 8 mg/kg intravenous infusion followed by 6 mg/kg intravenous infusion not more frequently than once every 3 weeks was given.^{7,8,9}

Guidelines

The use of trastuzumab products is recommended across multiple National Comprehensive Cancer Network (NCCN) guidelines and the NCCN Compendium.²⁻¹⁰ Recommendations with a category of evidence of 2B or higher support the approval criteria outlined below.

POLICY STATEMENT

Prior Authorization is recommended for medical benefit coverage of trastuzumab products. Approval is recommended for those who meet the Criteria and Dosing for the listed indications. Extended approvals are allowed if the patient continues to meet the **Criteria** and **Dosing**. Requests for doses outside of the established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with trastuzumab products, as well as the monitoring required for adverse events and long-term efficacy, approval requires trastuzumab products to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of trastuzumab products is recommended in those who meet one of the following:

FDA-Approved Indications

-
1. **Breast Cancer.** Approve for the duration noted if the patient meets ALL of the following (A, B, C, and D):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient has human epidermal growth factor receptor 2 (HER2)-positive disease; AND
 - C) Patient meets ONE of the following (i or ii):
 - i. Approve for 1 year (total) if trastuzumab is used for neoadjuvant (preoperative)/adjuvant therapy; OR
 - ii. Approve for 1 year if trastuzumab is used for recurrent or metastatic disease; AND
 - D) The medication is prescribed by or in consultation with an oncologist.

Dosing: Approve ONE of the following dosing regimens (A, B, or C):

- A) 4 mg/kg intravenously followed by 2 mg/kg not more frequently than once weekly; OR

- B) 8 mg/kg intravenously followed by 6 mg/kg not more frequently than once every 3 weeks; OR
- C) 4 mg/kg intravenously followed by 2 mg/kg not more frequently than once weekly during chemotherapy, then 6 mg/kg not more frequently than once every 3 weeks.

2. Gastric, Esophageal, or Gastroesophageal Junction Cancer. Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):

- A) Patient is ≥ 18 years of age; AND
- B) Patient has locally advanced or metastatic disease; AND
- C) Patient has human epidermal growth factor receptor 2 (HER2)-positive disease; AND
- D) Patient meets BOTH of the following (i and ii):
 - i. Trastuzumab will be used as first-line therapy; AND
 - ii. Trastuzumab will be used in combination with chemotherapy; AND

Note: Examples of chemotherapy are cisplatin, oxaliplatin, capecitabine, 5-fluorouracil (5-FU).
- E) The medication is prescribed by or in consultation with an oncologist.

Dosing. Approve ONE of the following dosing regimens (A or B):

- A) 8 mg/kg intravenously followed by 6 mg/kg not more frequently than once every 3 weeks; OR
- B) 6 mg/kg intravenously followed by 4 mg/kg not more frequently than once every 2 weeks.

Other Uses with Supportive Evidence

3. Appendiceal Cancer. Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):

- A) Patient is ≥ 18 years of age; AND
- B) Patient has human epidermal growth factor receptor 2 (HER2)-positive disease; AND
- C) Patient has not previously been treated with a HER2 antagonist; AND
- Note: Examples of HER2 antagonists include trastuzumab, Perjeta (pertuzumab intravenous infusion), or Phesgo (pertuzumab, trastuzumab, and hyaluronidase-zzxf subcutaneous injection).
- D) The medication is used in combination with Perjeta (pertuzumab intravenous infusion) or Tukysa (tucatinib tablets); AND
- E) The medication is prescribed by or in consultation with an oncologist.

Dosing: Approve 8 mg/kg intravenously followed by 6 mg/kg not more frequently than once every 3 weeks.

4. Biliary Tract Cancer. Approve for 1 year if the patient meets ALL of the following (A, B, C, D, E, and F):

- A) Patient is ≥ 18 years of age; AND
- B) Patient has unresectable or metastatic disease; AND
- C) Patient has human epidermal growth factor receptor 2 (HER2)-positive disease; AND
- D) The medication will be used in combination with Perjeta (pertuzumab intravenous infusion) or Tukysa (tucatinib tablets); AND
- E) The patient has tried one systemic regimen; AND
- Note: Examples of a systemic regimen include: gemcitabine and cisplatin, 5-fluorouracil and oxaliplatin, capecitabine and oxaliplatin, or gemcitabine and oxaliplatin.
- F) The medication is prescribed by or in consultation with an oncologist.

Dosing: Approve 8 mg/kg intravenously followed by 6 mg/kg not more frequently than once every 3 weeks.

-
- 5. Colon or Rectal Cancer.** Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):
- A) Patient is ≥ 18 years of age; AND
 - B) Patient has advanced or metastatic disease; AND
 - C) Patient has human epidermal growth factor receptor 2 (HER2)-positive disease; AND
 - D) The medication is used in combination with Perjeta (pertuzumab intravenous infusion) or Tukysa (tucatinib tablets); AND
 - E) The medication is prescribed by or in consultation with an oncologist.

Dosing: Approve ONE of the following dosing regimens (A or B):

- A) 8 mg/kg intravenously followed by 6 mg/kg not more frequently than once every 3 weeks; OR
- B) 4 mg/kg intravenously followed by 2 mg/kg not more frequently than weekly.

-
- 5. Endometrial Carcinoma.** Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):
- A) Patient is ≥ 18 years of age; AND
 - B) Patient has advanced or recurrent uterine serous carcinoma; AND
 - C) Patient has human epidermal growth factor receptor 2 (HER2)-positive disease; AND
 - D) Trastuzumab will be used in combination with chemotherapy; AND
Note: Examples of chemotherapy are carboplatin, paclitaxel.
 - E) The medication is prescribed by or in consultation with an oncologist.

Dosing: Approve 8 mg/kg intravenously followed by 6 mg/kg not more frequently than once every 3 weeks.

-
- 6. Salivary Gland Tumor.** Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):
- A) Patient is ≥ 18 years of age; AND
 - B) Patient has recurrent, unresectable, or metastatic disease; AND
 - C) Patient has human epidermal growth factor receptor 2 (HER2)-positive disease; AND
 - D) The medication is prescribed by or in consultation with an oncologist.

Dosing: Approve 8 mg/kg intravenously followed by 6 mg/kg not more frequently than once every 3 weeks.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of trastuzumab is not recommended in the following situations.

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Herceptin® intravenous infusion [prescribing information]. South San Francisco, CA: Genentech; May 2026.
2. The NCCN Breast Cancer Clinical Practice Guidelines in Oncology (version 4.2026 – June 12, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026.
3. The NCCN Colon Clinical Practice Guidelines in Oncology (version 2.2026 – April 7, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026.
4. The NCCN Rectal Clinical Practice Guidelines in Oncology (version 2.2026 – April 7, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026.
5. The NCCN Gastric Clinical Practice Guidelines in Oncology (version 3.2026 – June 3, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026.
6. The NCCN Esophageal and Esophagogastric Junction Cancers Clinical Practice Guidelines in Oncology (version 3.2026 – June 3, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026.
7. The NCCN Head and Neck Cancers Clinical Practice Guidelines in Oncology (version 2.2026 – May 12, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026.
8. The NCCN Biliary Tract Cancers Clinical Practice Guidelines in Oncology (version 1.2026– March 10, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026.
9. The NCCN Uterine Neoplasms Clinical Practice Guidelines in Oncology (version 3.2026– June 16, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026.
10. The NCCN Drugs & Biologics Compendium. © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026. Search term: trastuzumab.
11. Hercessi™ intravenous infusion [prescribing information]. Raleigh, NC: Accord BioPharma; September 2024.
12. The NCCN Appendiceal Neoplasms and Cancers Clinical Practice Guidelines in Oncology (version 2.2026 – April 10, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on June 19, 2026.

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	Biliary Tract Cancer: Added “Tukysa (tucatinib tablets)” as one of the agents that can be used in combination with trastuzumab.	07/17/2024
Selected Revision	Added Hercessi, a new biosimilar, to the policy.	12/11/2024
Update	04/21/2025: The policy name was changed from “Oncology (Injectable) - Trastuzumab Products UM Medical Policy” to “Oncology (Injectable - HER2 Antagonist) - Trastuzumab Products UM Medical Products Policy”	N/A
Annual Revision	No criteria changes	07/16/2025
Annual Revision	Appendiceal Cancer: Added new approval condition under “Other Uses with Supportive Evidence”. Colon or Rectal Cancer: Removed lapatinib from combination use with trastuzumab products.	06/24/2026

N/A – Not Applicable.